



(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 603 305 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

14.05.1997 Bulletin 1997/20

(21) Application number: **92919977.6**

(22) Date of filing: **08.09.1992**

(51) Int. Cl.⁶: **A61K 38/16**

(86) International application number:
PCT/US92/07556

(87) International publication number:
WO 93/05806 (01.04.1993 Gazette 1993/09)

(54) **USE OF A THYMOSIN IN THE TREATMENT OF HEPATITIS C**

VERWENDUNG EINES THYMOSINS BEI DER BEHANDLUNG VON HEPATITIS C

UTILISATION D'UNE THYMOSINE DANS LE TRAITEMENT DE L'HEPATITE C

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
SE**

(30) Priority: **13.09.1991 US 759544**

(43) Date of publication of application:
29.06.1994 Bulletin 1994/26

(73) Proprietors:

- **ALPHA 1 BIOMEDICALS, INC.**
Bethesda, MD 20817 (US)
- **THE BOARD OF GOVERNORS
OF WAYNE STATE UNIVERSITY**
Detroit Michigan 48202 (US)
- **THE UNITED STATES OF AMERICA**
as represented by the Secretary of the Army
Washington, D.C. (US)

(72) Inventors:

- **MUTCHNICK, Milton**
West Bloomfield, MI 48322 (US)

- **CHRETIAN, Paul**
Rockville, MD 20852 (US)
- **SHERMAN, Kenneth E.**
Aurora, CO 80015 (US)

(74) Representative: **Bunke, Holger, Dr.rer.nat. Dipl.-
Chem. et al**
Prinz & Partner
Manzingerweg 7
81241 München (DE)

(56) References cited:
EP-A- 0 331 934

- **CHEMICAL ABSTRACTS, vol. 106, no. 15, 13
April 1987, Columbus, Ohio, US; abstract no.
117943h, J.W. EICHBERG ET AL. 'EFFECT OF
THYMOSIN IMMUNOSTIMULATION WITH AND
WITHOUT CORTICOSTEROID
IMMUNOSUPPRESSION ON CHIMPANZEE
HEPATITIS B CARRIERS.' page 487-488 ;**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 603 305 B1

Description

This invention relates generally to a new use of a composition effective in the treatment of hepatitis C virus infection in patients.

Hepatitis C virus (HCV), the putative agent in the majority of post-transfusion acquired hepatitis, has been recently defined by a new serologic assay. Kuo, G., et al., *Science*, 244:362-4 (1989). Despite improvement in the quality of the blood-donor pool and the recent implementation of testing of donated blood, the current estimated incidence of acute infection among persons receiving transfusions is 5 to 10%. Alter, H.J., in Zuckerman, A.J., ed., *Viral Hepatitis and Liver Disease*, Allen K. Liss, New York, 1988, pp. 537-42. Chronic hepatitis develops in at least half the patients with acute HCV infection (representing about 90% of patients with non-A, non-B hepatitis (NANB)), and cirrhosis develops in at least 20% of this group. Thus, of the approximately 3 million persons who receive transfusions in the United States each year, acute hepatitis C will develop in about 150,000. Chronic hepatitis C will develop in at least 75,000 of these, and among them cirrhosis will develop in more than 15,000. Among patients with post-transfusion hepatitis, up to about 90% are positive for the HCV antibody. Davis, G.L., et al., *New England Journal of Medicine*, 321:1501-6 (1989). Patients with sporadic NANB hepatitis (no specific risk factors) are also very likely to have the anti-HCV antibody. Kuo, et al. (1989) above. While most of the patients who contract hepatitis C will have subclinical or mild disease, approximately 50% will progress to a chronic disease state characterized by fluctuating serum transaminase abnormalities and inflammatory lesions on liver biopsy. By some estimates, cirrhosis will develop in up to about 20% of this group. Koretz, R.L., et al., *Gastroenterology*, 88:1251-4 (1985).

With the aim of halting or slowing the progression of HCV-related diseases, a variety of drugs have been evaluated in recent years. Both acyclovir and corticosteroids (which are beneficial in autoimmune chronic active hepatitis) are ineffective. Pappas, S.C., *J. Med. Virol.*, 15:1-9 (1985); Stokes, P., et al., *Gastroenterology*, 92:1783 abstract (1987).

To date, α -interferon (IFA) appears to be the most promising candidate, although not necessarily the final answer. Hoofnagle, J.H., et al., in *Viral Hepatitis: 1981 International Symposium*, Philadelphia, Franklin Institute Press, 1982, pp. 573-83; Hoofnagle, J.H., et al., *New England Journal of Medicine*, 315:1575-8 (1986); Thomson, J., *Lancet*, 1:539-41 (1987); Kiyosawa, K., et al., in Zuckerman, A., ed., *Viral Hepatitis and Liver Disease*, Allen K. Liss, New York, 1983, pp. 895-7. Hoofnagle, J.H., et al., *Sem. Liver Dis.*, 9:259-263 (1985). The interferons are host proteins made in response to viral infections as well as other antigenic stimuli. They are classified by their cell of origin as well as their antigenicity. α -Interferon is made by lymphoblastoid cells, β -interferon by fibroblasts, and γ -interferon by T-cells. Subtypes in each group are based on antigenic/structural characteristics. Recombinant forms for each group have been developed and are commercially available. A pilot study utilizing IFA on ten patients with well-characterized post-transfusion NANB hepatitis was reported in 1986 by Hoofnagle et al. (Hoofnagle, J.H., et al., *New England Journal of Medicine*, 315:1575-8 (1986)). In this study, eight of ten patients improved their serum alanine transaminase (ALT) levels within one month of starting therapy. IFA therapy consisted of 5 million units (MU) daily in seven of the patients and one MU daily in three patients. In all subjects the dose was gradually reduced to 1 MU daily and then finally switched to an alternate day or every three day regimen. In three patients who had post-treatment liver biopsies, the specimen showed a marked improvement in the degree of portal inflammation and loss of parenchymal hepatocytic necrosis. Side effects were common at the 5 MU/day dose and virtually absent at 1 MU/day.

The effects of recombinant human interferon α in a prospective, randomized, double-blind, placebo-controlled trial in patients with well-documented chronic HCV infection has recently been carried out. Di Bisceglie, A.M., et al., *New England Journal of Medicine*, 321:1506-10 (1989). Forty-one patients were enrolled in the trial, 37 of whom were later found to have antibody to HCV. Twenty-one patients received interferon α (2 MU) subcutaneously three times weekly for six months, and twenty received placebo. The mean serum ALT and the histological features of the liver improved significantly in the patients treated with interferon, but not in the patients given placebo. Ten patients treated with interferon (48%) had a complete response, defined as a decline in mean serum ALT to the normal range during therapy; three others had a decrease in mean ALT of more than 50%. After treatment ended, however, serum ALT usually returned to pretreatment levels; six to twelve months after the discontinuation of interferon therapy, only two patients (10%) still had normal values. The authors concluded that interferon α therapy is beneficial in reducing disease activity in chronic hepatitis C; however, the beneficial responses are often transient and side effects are known to appear.

In another, broader study, chronic hepatitis C (NANB hepatitis) in 166 patients was treated with either 3 MU or 1 MU of recombinant human α -IFA three times weekly for 24 weeks or to no treatment. The serum ALT level became completely normal in 22 of the 26 patients (85%) who responded to treatment with 3 MU of interferon, and nine of the sixteen patients (56%) responded to treatment with 1 MU. The patients who received 3 MU of interferon had histologic improvement because of the regression of lobular and periportal inflammation. However, relapse within six months after the completion of treatment occurred in 51% of the patients treated with 3 MU of interferon and in 44% of those treated with 1 MU. Davis, G.L., et al., *New England Journal of Medicine*, 321:1501-06 (1989). These authors concluded that a 24-week course of interferon therapy is effective in controlling disease activity in many patients with hepatitis C, although relapse after the cessation of treatment is common.

A multi-center randomized control trial of recombinant human α -IFN in patients with chronic NANB hepatitis has

been reported recently. Marcellin, P., et al., Hepatology, 13:393-97 (1991). Patients were randomly assigned to no treatment or to 1 to 3 MU of α -interferon given three times a week for 24 weeks. Forty-five patients (75%) were positive for antibody to HCV. During the 24 week treatment period, mean serum ALT levels decreased in both treatment groups, but the decrease was statistically significant only in the 3 MU group. However, at 24 weeks, the proportion of patients with normal ALT levels was similar in the 3 MU group (39%) and the 1 MU group (45%) and both were significantly higher than in controls (0%). Repeat liver biopsy specimens showed a significant decrease in the severity of histological changes in the higher dose group but not in the lower dose group or in controls. However, after treatment, the mean ALT levels rose in both treated groups. The proportion of patients with normal ALT levels at week 48 was 28% in the 3 MU group and 20% in the 1 MU group. The authors conclude that a dose of 3 MU was superior to 1 MU of α -interferon given three times per week for 24 weeks in inducing improvements in serum ALT levels and liver histological examinations. However, relapse in disease activity occurred in approximately half of the responders when interferon was stopped. The response to α -interferon did not correlate with the source of infection or with the presence or absence of anti-HCV antibody titres in patient sera.

It is clear, therefore, that while α -interferon has a beneficial effect on the course of HCV infection, this effect is frequently only transient. Therefore, new modalities are necessary in order permanently to eradicate the effects of hepatitis C virus on the patient.

Another class of polypeptide immune modifiers derived from the thymus gland, the thymosins, has been shown to trigger maturational events in lymphocytes, to augment T-cell function and to promote reconstitution of immune defects. Low, T.L.K., et al., "Thymosins: Structure, Function and Therapeutic Application", Thymus, 6:27-42 (1984).

Thymosin Fraction Five (TF-5), originally described by Goldstein et al. (Proc. Nat'l Acad. Sci. (USA), 69:1800-1803 (1972)), is a partially purified extract of bovine thymus containing at least 40 peptide components, 20 of which have been purified to homogeneity or near homogeneity; it contains about 0.6% of Thymosin α -1 (THN α ₁). Low, 1984, above.

THN α ₁, initially isolated from TF-5, has been sequenced and chemically synthesized. Wetzel, R., et al., Biochemistry, 19:6096-6104 (1980). Its sequence is highly homologous in mice, calves and humans. THN α ₁ is a 28 amino acid polypeptide with a molecular weight of 3100 that has shown activity qualitatively similar to TF-5 in modulating the immune system. Low, T.L.K., et al., J. Biol. Chem., 254:981-6 (1979). THN α ₁ has potent immunologic activity, including stimulation of α - and γ -interferon production, increasing macrophage migration inhibitory factor production, inducing expression of T-cell markers, including IL-2 receptors, and improving T-cell helper cell activity. Schulof, R.S., et al., in The Lymphocyte, Allen J. Liss Inc., New York, 1981, pp. 191-215; Low, T.L.K., et al., in "Thymosins: Structure, Function and Therapeutic Applications", Thymus, 6:27-43 (1984); Koutab, N.M., et al., Immunopharm., 16:97-105 (1988). Studies in mice have demonstrated a synergistic effect of THN α ₁ and interferon on natural killer-cell activity in immunosuppressed mice. Favilli, C., et al., Cancer Immunol. Immunother., 20:189-92 (1985). TF-5 and THN α ₁ can influence immunoregulatory T-cell function, promote production of interferon- α , interferon- γ and interleukin-2 by human lymphocytes and increase interleukin-2 receptor expression. Marshall, G.D., et al., J. Immunol., 126:741-4 (1981); Mutchnick, M.G., et al., Clin. Immunol. Immunopathol., 23:626-33 (1982); Szein, M.B., et al., Proc. Nat'l Acad. Sci. (USA), 83:6107-6111 (1986); Serrate, S.A., et al., J. Immunol., 1939:2338-43 (1987); Baxevis, C.N., et al., Immunopharm., 13:133-41 (1987); and, Svedersky, L.P., Eur. J. Immunol., 12:244-7 (1982).

Clinical trials of TF-5 and THN α ₁ as primary or adjunctive therapy in patients with immunodeficiency or cancer indicate that these agents enhance immune responsiveness and augment specific lymphocyte functions. Clinical trials of TF-5 and purified THN α ₁ have been underway for a number of years. Early trials in patients with cancer or immunodeficiency states were encouraging, though not definitive. Goldstein, A.L., et al., Transp. Proc., 9:1141 (1977); Barrett, D.J., et al., J. Pediatr., 97:61 (1980); and Cohen, M.H., et al., J. Amer. Med. Assoc., 241:1813-5 (1979). THN α ₁ use has been described in a randomized trial of patients with nonsmall cell lung cancer. Patients were treated with THN α ₁ at a dose of 900 μ grams/m² subcutaneously twice weekly or daily for two weeks and then twice weekly after completing a course of radiotherapy. The only side effect of THN α ₁ was mild burning at the injection site in three patients. This was attributed to the drug lot and may have been due to the carrier preparation. Relapse-free survival and overall survival were greater in both THN α ₁ treatment groups than in the placebo group; some restoration of radiation-suppressed immune function was also noticed. There was no increase in T-cell numbers associated with this. Schulof, R.S., et al. J. Biol. Response Modifiers, 4:147-58 (1985).

Recent double-blind, randomized trials with thymosins have been performed in elderly men in an effort to increase response to influenza vaccine. Gravenstein, S., et al., JAGS, 37:1-8 (1989). Patients received synthetic THN α ₁ subcutaneously twice weekly starting at the time the influenza vaccine was given. At six weeks post-vaccine, those patients randomized to receive the drug had higher levels of antibody to influenza than controls. This difference was accentuated in the very elderly (ages 77-99). No clinical or biochemical toxicity was observed in drug recipients.

There are preliminary reports that thymosins may be effective against infections caused by hepatitis viruses other than HCV. In an animal model of viral hepatitis, the woodchuck infected with the Woodchuck Hepatitis Virus, THN α ₁ suppressed viral DNA replication, but produced no improvement in clinical parameters. Korba, B.E., et al., Hepatology, 12:Abs. 880 (1990). In a pilot clinical trial with patients with Chronic Active Hepatitis B caused by the hepatitis B virus

(HBV), patients treated for a year with THN α_1 (5 patients) or with TF-5 (2 patients) showed a marked decrease in serum ALT; 6 of the 7 patients also showed reduced levels of serum HBV DNA, and 5 of 6 patients initially positive for serum hepatitis B surface antigen (HBsAg) subsequently cleared this antigen. Mutchnick, M.C., et al., Hepatology, 10:Abs. 575 (1989). No suggestion was made in these abstracts that the thymosins would be effective against any other hepatitis viruses.

There remains, therefore, an important need in the art for a new modality for the treatment of HCV infections in mammals.

EP-A1-0 331 934 discloses a pharmaceutical composition containing thymosin α_1 , human albumin as a stabilizer and, optionally, an aqueous pharmaceutically acceptable vehicle. There are several medical indications of said pharmaceutical composition given in said document: Protection from undesirable effects of the immunosuppressive agents employed in cancer therapy, preventive activity against progression of leukemias in mice, and protective activity against opportunistic infections such as infections with *Candida*, *Pseudomonas* and *Serratia* strains.

A treatment modality for HCV infections has been devised comprising the administration to mammals of immune system-potentiating doses of one or more thymosins, alone or in combination with interferon therapy.

It is thus an object of this invention to provide a novel medical indication of compositions in the treatment of mammals with one or more members of the thymosin family of biological response modifiers.

It is another object of this invention to provide a further medical indication of compositions in the treatment of mammals comprising combination therapy with one or more thymosins and one or more interferons.

The present invention resides in the use of a composition comprising an immune system-potentiating amount of at least one thymosin or an immune system-potentiating thymosin fragment for the manufacture of a medicament for the treatment of a mammal infected with hepatitis C virus.

According to preferred embodiments of the present invention, said thymosin is selected from the group consisting of Thymosin Fraction Five and Thymosin α_1 , and said composition further comprises an anti-viral effective amount of at least one interferon.

Other preferred embodiments of the present invention are defined in the subclaims appended.

Optionally, said composition used according to the present invention further comprises a pharmaceutically acceptable carrier.

Based on the present invention, a novel modality for treating HCV infection in mammals has been devised, comprising the administration to such mammals of one or more thymosins at doses which potentiate immune responses, alone or in combination with anti-viral doses of one or more interferons.

By the term "thymosins" is meant any or all of the immune system potentiating polypeptides naturally occurring in the thymus gland or produced by chemical or recombinant means, or fragments derived from any of these polypeptides. By the term "mammals" is meant any mammalian subject, including human and animal patients, requiring treatment for hepatitis C infection. "Mammal" and "subject" are used interchangeably.

Thymosin preparations suitable for treating HCV infections include TF-5, THN α_1 and fragments thereof, e.g., C-terminal 4-28 and 15-28, and N-terminal 1-8, 1-14 and 1-20 fragments. These may be obtained from Alpha-1 Biomedicals Inc., Foster City, California.

Subjects, e.g., human patients, may receive the thymosin by subcutaneous injection or infusion, at appropriate intervals for an appropriate period of time. The thymosin is administered to mammals infected with hepatitis C virus in amounts which facilitate or promote *in vivo* inactivation of hepatitis C virus. A pharmaceutical dosage unit of an immune system-potentiating amount of a thymosin, such as TF-5, can be from about 900 to about 1200 mg/m² body surface area in a pharmaceutically acceptable carrier. A pharmaceutical dosage unit of an immune system-potentiating amount of a thymosin, such as THN α_1 or immune system-potentiating fragments thereof, can be from about 900 to about 1200 μ g/m² body surface area in a pharmaceutically-acceptable carrier. Lyophilized preparations of thymosins or fragments which contain mannitol and phosphate buffer are dissolved in diluent prior to dispensing. Thymosins in diluent should remain stable for at least six months when stored in a refrigerator, it is convenient to dispense thymosin solutions in one ml dose vials. Some patients may require about eight vials per month.

For a typical human patient, an administration regimen of twice weekly (e.g., Monday and Thursday) subcutaneous injection of about 1500 to about 1700 μ g of THN α_1 or fragments therefrom is convenient. Dosages and length of treatment can be flexible, and can be determined by the subject's clinical response to the thymosins.

The course of the disease and its response to drug treatments may be followed by clinical examination and laboratory findings. As elevated serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) are known to occur in uncontrolled hepatitis C, and as a complete response to treatment is generally defined as the normalization of these serum enzymes, particularly ALT (Davis, G.L., et al., New England Journal of Medicine, 321:1501-6 (1989)), progress of treatment with thymosins is conveniently followed by this art-recognized test performed, e.g., on a sequential multiple analyzer.

Another means of evaluating subjects having antibodies to HCV (not all subjects with hepatitis C have detectable antibody to HCV - Weiner, A.J., et al., Lancet, 335:1-3 (1990)) is to periodically test subjects' sera for the titer of these antibodies. Anti-HCV antibodies may be tested by the currently available C 100-3 test (Kuo, G., et al., Science, 244:362-

4 (1989)), by an Elisa test (Ortho Diagnostic Systems, Raritan, N.J.) or by a recombinant assay (RIBA-1 and RIBA-2, Chiron Corporation, Emeryville, CA). Any suitable test may be used.

In order to follow the course of HCV replication in subjects in response to drug treatment, HCV RNA may be measured in serum samples by, for example, a nested polymerase chain reaction assay that uses two sets of primers derived from the NS3 and NS4 non-structural gene regions of the HCV genome. Farci, P., et al., New England Journal of Medicine, 325:98-104 (1991); Ulrich, P.P., et al., J. Clin. Invest., 86:1609-14 (1990).

Other appropriate laboratory tests to follow the course of treatment are listed in Example 1 below.

Thymosin therapy may also be used in combination with interferon therapy, thereby combining the immune system potentiating effect of thymosins with the anti-viral effects of the interferons. An improved response rate at the currently used interferon doses would be beneficial, particularly in the light of dose-limiting side effects at higher doses of these proteins. An offshoot of this concept is the ability to achieve comparable efficacy with interferon plus thymosin at lower doses than would be required with interferon alone.

In this combination therapy regimen, one or more interferons (for example, recombinant interferon α -2b, Intron-A, Schering-Plough, Kenilworth, New Jersey) is (are) administered subcutaneously to subjects, e.g., human patients, at doses ranging between about 1 MU and 3 MU along with or sequentially with one or more thymosins, preferably including THN α_1 , at a dose of about 900 to about 1200 $\mu\text{g}/\text{m}^2$ body surface area.

Although the example above speaks in terms of recombinant interferon α -2b, other anti-HCV-effective interferons such as α -, β - and γ -interferons, recombinant or naturally occurring, may be advantageously used in this invention.

This combination dose regimen is flexible, and depends on the clinical condition of the subject. Where subjects are refractory to the preferred dosage levels, these may be increased within the limits dictated by undesirable side effects. Typically, injections are made five times per week and continue until an acceptable response by the subject is realized.

Tests to determine the effectiveness of the combination therapy may be the same as those described above for thymosin treatment alone. In addition, histological examination of liver biopsy samples may be used as a second major criteria for evaluation. Knodell, R.G., et al., Hepatology, 1:431-5 (1981), whose Histological Activity Index (portal inflammation, piecemeal or bridging necrosis, lobular injury and fibrosis) provides a scoring method for disease activity.

The following examples are provided merely to illustrate the invention, and are not to be construed in any way as limiting the scope of invention as set forth in the specification and claims.

EXAMPLE 1

Preparation of Injectable Formulation

Pharmaceutical dosage units of 1 ml each are prepared from the ingredients shown in Table 1 below.

TABLE 1

Active Ingredient	Amount Per mL
Thymosin α -1	0.0016 g
Inactive Ingredients	
mannitol, U.S.P.	0.050 g
sodium phosphate dibasic, heptahydrate, U.S.P.	0.002 g
sodium phosphate monobasic, monohydrate, U.S.P.	0.0005 g
sodium phosphate dibasic, 2 mg/ml solution	
sodium phosphate monobasic, 0.5 mg/ml solution	
water for injection, U.S.P.	

EXAMPLE 2

Treatment of Hepatitis C Infections in Human Patients with Thymosins and Interferons

Adult patients with chronic active hepatitis C (CAHC) are randomized to one of four study groups, made up of about 40 patients per group. Selection criteria include: (1) patients are adults (at least 18 years of age); (2) serum ALT is ele-

vated for at least six months prior to treatment with at least one value greater than twice the upper limit of normal in the laboratory doing the testing; (3) patients test positive for HCV antibody on two occasions and on a confirmatory test; and (4) liver biopsy within three months of treatment exhibits pathology consistent with chronic active hepatitis.

Exclusion criteria include: (1) recent use of other anti-viral or immunosuppressive medication; (2) hemophilia, pregnancy or HIV infection, or other serious illness that could prevent completion of the course of treatment; (3) other forms of liver disease, including hepatitis A or B, α -1 antitrypsin deficiency, Wilson's disease, and hemochromatosis must be absent; (4) autoimmune markers (ANA, ASMA, AMA, anti-LKMI) must be absent or, if present, titers should be $< 1:40$; (5) leukocyte deficiency ($< 3,000$); (6) low absolute neutrophil count ($< 1,000$); (7) low platelets ($< 75,000$); (8) low Hb (< 11 g/dL); (9) high bilirubin (> 4 mg/dL); and (10) low serum albumin (3 g/dL).

The first of the four randomized groups receives interferon, preferably interferon α -2b, at a dose of 3 million units (MU) subcutaneously (SQ) on Mondays, Wednesdays and Fridays, and receives placebos on Tuesdays and Saturdays. The second group receives the same dose/schedule of interferon, plus a thymosin, preferably THN α_1 , at a dose of 900 μ g/m² SQ on Tuesdays and Saturdays. The third group receives the same dose/schedule of a thymosin alone. The fourth group receives placebo treatment initially, but can be randomized to the three treatment groups thereafter. Interferons and thymosins can be recombinant.

Patients begin treatment while hospitalized for about one week, during which period side-effects are monitored.

Outpatient follow-up is initially at one week intervals for two weeks, then at two week intervals for two months, and then monthly for the remainder of the treatment period. At each visit the following lab tests are performed: CBC, platelet count, differential and ESR, ALT, AST, GGT, alkaline phosphatase, bilirubin, total bilirubin/albumin and HCV antibody.

At monthly intervals serum γ -globulin, TSH, ANA and ASMA are assessed.

Drug toxicity is monitored on an ongoing basis using both clinical and laboratory parameters.

Within one month of completing the initial six months of treatment, patients undergo liver biopsy for pathological examination according to Knodell et al. above. This system provides a numerical scoring system of histological activity in patients with asymptomatic CAH.

At this time, control patients are randomized into three groups to receive one of the three treatment modalities, assuming that they still have CAH on follow-up liver biopsy, and that one arm of the study does not show highly significant positive or negative results on analysis at six months.

Patients in the treatment groups are followed to evaluate recrudescence of disease as evidenced by rising ALT levels. Patients who showed response in the initial six month treatment period, but who have a recurrence of the disease thereafter, are provided with additional therapy.

Additional serum or tissue tests are performed if possible: evaluation of antibodies to interferons and thymosins, polymerase chain reaction amplification of hepatitis C genome segments in liver biopsy samples, and quantitative evaluation of anti-hepatitis C serum titers.

EXAMPLE 3

The treatment protocol is as in Example 2, except that the interferon is used at the level of 2 MU, and the thymosin at 1050 μ g/m².

EXAMPLE 4

The treatment protocol is as in Example 3, except that 1 MU of the interferon and 1200 μ g/m² of the thymosin are used.

EXAMPLE 5

Analysis of Data

There are two primary criteria for response to therapy - normalization of ALT levels by the end of the treatment period (a partial response may be defined as a decrease of at least 50% of initial ALT), and histological improvement as determined by the Histological Activity Index (HAI) of Knodell et al. above.

This analysis provides a raw score ranging from 1 to 22 per sample. Paired data can be analyzed using the Wilcoxon paired-sample test. Additionally, samples can be classified into mild, moderate or reverse CAH, and improvement assessed using the Chi-square statistical analysis.

Life-table analysis is used to evaluate remission and relapse status in terms of normalization of ALT levels. Other continuous variables are analyzed using Student's t test. Dichotomous data are subjected to Chi square or Fisher's exact test, as is appropriate.

A power analysis was done to determine the number of patients in each test group in order to show predicted differences. Power analysis applied to an ANOVA using a power of 0.80 with $\alpha = 0.05$, coupled with prior studies of mean

ALT levels and their variances, estimated a need for 21 to 52 patients in each test group to show a mean ALT difference of 15 IU/L. As 3 to 5% of patients are expected to drop out, and factoring in treatment of the control group after six months, 40 patients per group was arrived at.

5 Claims

1. Use of a composition comprising an immune system-potentiating amount of at least one thymosin or an immune system-potentiating thymosin fragment *for the manufacture of a medicament for the treatment of a mammal infected with hepatitis C virus.*
2. The use of claim 1, wherein said thymosin is selected from the group consisting of Thymosin Fraction Five and Thymosin α -1.
3. The use of claim 1 further comprising an anti-viral effective amount of at least one interferon.
4. The use of claim 3, wherein said interferon is α -interferon and wherein said anti-viral effective amount is from about 1 million to about 3 million units of said interferon.
5. The use of claim 3, wherein said interferon is selected from the group consisting of α -, β -, and γ -interferons.
6. The use of claim 5, wherein said α -interferon is interferon α -2B.
7. The use of claim 6, wherein said interferon is recombinant interferon.
8. The use of claim 1, wherein said thymosin is Thymosin α -1, said immune system-potentiating amount is a human immune system-potentiating amount, and said amount is from about 900 to about 1200 $\mu\text{g}/\text{m}^2$ body surface area of said human.
9. The use of claim 1, wherein said thymosin is Thymosin Fraction Five, said immune system-potentiating amount is a human immune system-potentiating amount, and said amount is about 167 times the amount of Thymosin α -1, per m^2 body surface area of said human.
10. The use of claim 1, wherein said thymosin is Thymosin α -1, and wherein said amount is about 1500 to about 1700 μg of Thymosin α -1, per pharmaceutical dosage unit.
11. The use of any one of claims 1 to 10, wherein said composition further comprises a pharmaceutically acceptable carrier.

Patentansprüche

1. Verwendung einer Zusammensetzung, die eine das Immunsystem verstärkende Menge an mindestens einem Thymosin oder einem das Immunsystem verstärkenden Thymosinfragment umfaßt, zur Herstellung eines Arzneimittels zur Behandlung eines mit dem Hepatitis C-Virus infizierten Säugers (Mensch und Tier).
2. Verwendung nach Anspruch 1, bei der das Thymosin ausgewählt wird aus der aus der Thymosinfraktion Fünf und Thymosin α -1 bestehenden Gruppe.
3. Verwendung nach Anspruch 1, die zusätzlich eine antiviral wirksame Menge an mindestens einem Interferon umfaßt.
4. Verwendung nach Anspruch 3, wobei das Interferon α -Interferon ist und wobei die antiviral wirksame Menge etwa 1 Million bis etwa 3 Millionen Einheiten des Interferons beträgt.
5. Verwendung nach Anspruch 3, wobei das Interferon ausgewählt wird aus der aus α -, β - und γ -Interferonen bestehenden Gruppe.
6. Verwendung nach Anspruch 5, bei der das α -Interferon Interferon α -2B ist.
7. Verwendung nach Anspruch 6, bei der das Interferon rekombinantes Interferon ist.

8. Verwendung nach Anspruch 1, wobei das Thymosin Thymosin α -1 ist, die das Immunsystem verstärkende Menge eine das Immunsystem des Menschen verstärkende Menge ist und diese Menge etwa 900 bis etwa 1200 $\mu\text{g}/\text{m}^2$ Körperoberfläche des Menschen beträgt.
- 5 9. Verwendung nach Anspruch 1, wobei das Thymosin Thymosinfraktion Fünf ist, die das Immunsystem verstärkende Menge eine das Immunsystem des Menschen verstärkende Menge ist und diese Menge etwa das 167fache der Menge an Thymosin α -1 pro m^2 Körperoberfläche des Menschen ist.
- 10 10. Verwendung nach Anspruch 1, wobei das Thymosin Thymosin α -1 ist und die Menge etwa 1500 bis etwa 1700 μg Thymosin α -1 pro pharmazeutischer Dosierungseinheit beträgt.
11. Verwendung nach einem der Ansprüche 1 bis 10, bei der die Zusammensetzung zusätzlich einen pharmazeutisch verträglichen Träger umfaßt.

15 Revendications

1. Utilisation d'une composition comprenant une quantité potentialisatrice du système immunitaire, d'au moins une thymosine, ou d'un fragment de thymosine, potentialisateur du système immunitaire, pour la fabrication d'un médicament pour le traitement d'un mammifère contaminé par le virus de l'hépatite C.
- 20 2. Utilisation selon la revendication 1, dans laquelle ladite thymosine est choisie au sein du groupe comprenant la Fraction Cinq de Thymosine, et la thymosine α -1.
3. Utilisation selon la revendication 1, comprenant aussi une quantité antivirale efficace d'au moins un interféron.
- 25 4. Utilisation selon la revendication 3, dans laquelle ledit interféron est l'Interféron- α , et dans laquelle ladite quantité antivirale efficace est comprise entre environ 1 million à environ 3 millions d'unités dudit interféron.
5. Utilisation selon la revendication 3, dans laquelle ledit interféron est choisi au sein du groupe comprenant les Interférons- α - β , et γ .
- 30 6. Utilisation selon la revendication 5, dans laquelle ledit Interféron α est l'Interféron α -2B.
7. Utilisation selon la revendication 6, dans laquelle ledit interféron est de l'interféron recombinant.
- 35 8. Utilisation selon la revendication 1, dans laquelle ladite thymosine est de la Thymosine α -1, ladite quantité potentialisatrice du système immunitaire est une quantité potentialisatrice du système immunitaire humain, et ladite substance est comprise entre environ 900 et environ 1.200 μg par mètre carré de surface corporelle dudit humain.
- 40 9. Utilisation selon la revendication 1, dans laquelle ladite thymosine est la Fraction Cinq de Thymosine, ladite quantité potentialisatrice du système immunitaire est une quantité potentialisatrice du système immunitaire humain, et ladite quantité est environ 167 fois celle de Thymosine α -1, par mètre carré de surface corporelle dudit humain.
- 45 10. Utilisation selon la revendication 1, dans laquelle ladite thymosine est de la Thymosine α -1, et dans laquelle ladite quantité est comprise entre environ 1.500 et environ 1.700 μg de Thymosine α -1, par dose pharmaceutique unitaire.
- 50 11. Utilisation selon l'une quelconque des revendications 1 à 10, dans laquelle ladite composition contient aussi un véhicule pharmaceutiquement acceptable.